

cells, although the exact nature of the genetic events responsible for this process could not be identified by these techniques. As such, the potentially invasive nature of this tumor should be emphasized, and adequate, complete excision should be performed at the time of presentation.

We gratefully acknowledge the assistance of Maureen Parker and Almaz Isaac for assistance with tumor sample preparation and Michael Grande for the preparation of the fibroblast and tumor DNA samples. The PGK probes were kind gifts of Dr. Katherine Siminovitch.

This research was supported by NIH grant AG-09927 (MP) and the Dermatology Foundation (HCT).

REFERENCES

1. Knudsen AG Jr: Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 68:820-823, 1971
2. Vogelstein B, Fearon ER, Hamilton SR, Preisinger AC, Willard HF, Michelson AM, Riggs AR, Orkin SH: Clonal analysis using recombinant DNA probes from the X-chromosome. *Cancer Res* 47:4806-4813, 1987
3. Gartler SM, Ziprowski L, Krakowski A, Ezra R, Szeinberg A, Adams A: Glucose 6-phosphate dehydrogenase mosaicism as a tracer in the study of hereditary multiple trichoepithelioma. *Am J Hum Genet* 18:282-287, 1966
4. Fialkow PJ, Sagabiel RW, Gartler SM, Rinoio D: Multiple cell origin of hereditary neurofibromas. *N Engl J Med* 248:298-300, 1971
5. Friedman JM, Fialkow PJ, Greene CL, Weinberg MN: Probable clonal origin of neurofibrosarcoma in a patient with hereditary neurofibromatosis. *J Natl Cancer Inst* 69:1289-1291, 1982
6. Laskin WB: Dermatofibrosarcoma protuberans. *CA* 42:116-123, 1992
7. Bridge AG, Neff JR, Sandberg AA: Cytogenetic analysis of dermatofibrosarcoma protuberans. *Cancer Genet Cytogenet* 49:199-202, 1990
8. Lyon MF: X-chromosome inactivation and developmental patterns in mammals. *Biol Rev* 47:1-35, 1972
9. Botstein D, White RL, Skolnick M, Davis R: Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am J Hum Genet* 32:314-331, 1980
10. Keith DH, Singer-Sam J, Riggs AD: Active X chromosome DNA is unmethylated at eight CCGG sites clustered in a guanine-plus-cytosine-rich island at the 5' end of the gene for phosphoglycerate kinase. *Mol Cell Biol* 6:4122-4125, 1986
11. Peacocke M, Siminovitch KA: Linkage of the Wiskott-Aldrich syndrome with polymorphic DNA sequences from the human X-chromosome. *Proc Natl Acad Sci USA* 84:3430-3433, 1987
12. Feinberg AP, Vogelstein B: A technique for radiolabeling DNA restriction fragments to high specific activity. *Anal Biochem* 132:6-13, 1983
13. Fidler IJ, Talmadge JE: Evidence that intravenously derived murine pulmonary metastases can originate from the expansion of a single tumor cell. *Cancer Res* 46:5167-5171, 1986
14. Sidransky D, Frost P, Von Eschenbacher A, Oyasu R, Preisinger AC, Vogelstein B: Clonal origin of bladder cancer. *N Engl J Med* 326:737-740, 1992

ANNOUNCEMENT

The 42nd annual Symposium on the Biology of Skin will be "Genetics of Skin Disease." The symposium will be held at Snowmass Lodge in Snowmass, Colorado on July 24-28, 1993. The meeting will be held just before the annual meeting of the Society for Pediatric Dermatology. Those wishing to present a poster at the SBS may submit an abstract to the Symposium Director on a standard SID or ESDR abstract form. All correspondence regarding participation in this meeting, and all abstract submissions, should be addressed to: David A. Norris, M.D., Symposium on the Biology of Skin, Department of Dermatology B-144, University of Colorado School of Medicine, 4200 East Ninth Avenue, Denver, Colorado 80262.